

Certificate of Analysis



Recombinant Human FGF-basic 147
Catalog Number: 100-28
Lot Number: 0313-100-28

Product Description

Alternate Names: FGF2, HBGF-2, Prostatropin **Accession Number:** P09038

Description: Basic fibroblast growth factor (bFGF), also known as FGF2, is expressed by endothelial cells and is a mediator of angiogenesis. bFGF also has cardioprotective functions during heart injury. The application of bFGF is a critical component for embryonic stem cell culture systems and is necessary for maintaining cells in an undifferentiated state. Degredation of the full length bFGF N-terminus results in a truncated bFGF147aa protein when the protein is isolated from biological sources. The N-terminus extensions influence the localization of bFGF within the cell but do not affect the biological activity of bFGF. Therefore, there are no detectable differences in biological activity between the full length bFGF-154aa and the truncated bFGF-147aa recombinant proteins.

Source: Genetically modified *E.coli*.

Predicted MW: Monomer, 16.5 kDa (147 amino acids)

AA Sequence: MPALPEDGGS GAFPPGHFKD PKRLYCKNGG FFLRIHPDGR VDGVREKSDP HIKLQLQAE E RGVVSIK GVC ANRYLAMKED GRLLASKCVT DECFFFERLE SNNYNTYRSR KYTSWYVALK RTGQYKLGSK TGPGQKAILF LPMSAKS

Formulation: Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 75 mM sodium chloride, pH 7.5

Product Specifications

Specification	Method of Determination	Acceptance Criteria	Result
Purity	Reducing and Non-reducing SDS PAGE	≥95%	>95%
Endotoxin	Kinetic LAL (50% confidence)	≤ 1 EUs/ug	<0.05 EUs/ug
Biological Activity	BALB/c 3T3 cell proliferation	ED50 ≤5 ng/mL (≥2.0 x 10 ⁵ units/mg)	ED50 = 0.13 ng/mL (7.7 x 10 ⁶ units/mg)

Product Preparation and Storage

Country of Origin: USA **Shipping:** Room temperature **Storage Prior to Reconstitution:** -20°C

Reconstitution: Sterile water at 0.1 mg/mL **Date of Recon.:** _____ **Initials:** _____

Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialled to compensate for this loss.

Expiration Date: _____ **Date Received:** _____

12 months from date of receipt when stored at -20°C to -80°C as supplied.

1 month when stored at 4°C after reconstituting as directed.

3 months when stored at -20°C to -80°C after reconstituting as directed.

THIS PRODUCT IS FOR RESEARCH USE ONLY AND IS NOT FOR USE IN HUMANS!

Stephan E. M. 10/5/16
Quality Control