

Product Description

Alternate Names: FGF2, HBGF-2, Prostatropin **Accession/Uniprot Number:** P13109

Description: Basic fibroblast growth factor (FGF-basic), also known as FGF-2, is expressed by endothelial cells and is a mediator of angiogenesis. FGF-basic also has cardioprotective functions during heart injury. FGF-basic binds heparin in order to signal through fibroblast growth factor (FGF) receptor tyrosine kinases.

Source: Genetically modified *E.coli*. **This product is produced with no animal or human origin raw products. All processing and handling employs animal free equipment and animal free protocols.**

Predicted MW: Monomer, 16.4 kDa (146 amino acids)

AA Sequence: MPALPEDGGG AFPPGHFKDP KRLYCKNGGF FLRIHPDGRV DGVREKSDPH VKLQLQAEER GVVSIKGVCA NRYLAMKEDG RLLASKCVTE ECFFFERLES NNYNTYRSRK YSSWYVALKR TGQYKLGSKT GPGQKAILFL PMSAKS

Formulation: Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5

Product Specifications

Specification	Method of Determination	Acceptance Criteria	Result
Purity	Reducing and Non-reducing SDS-PAGE	≥95%	>95%
Endotoxin	Kinetic LAL (50% confidence)	≤ 1 EUs/ug	<0.05 EUs/ug
Biological Activity	NR6R-3T3 proliferation	ED50 ≤1 ng/mL (≥1.0 x 10 ⁶ units/mg)	ED50 = 0.0903 ng/mL (1.1 x 10 ⁷ units/mg)

Product Preparation and Storage

Country of Origin: USA **Shipping:** Room temperature **Storage Prior to Reconstitution:** -20°C

Reconstitution: Sterile water at 0.1 mg/mL **Date of Recon.:** _____ **Initials:** _____

Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws.

Expiration Date: _____ **Date Received:** _____

12 months from date of receipt when stored at -20°C to -80°C as supplied.

1 month when stored at 4°C after reconstituting as directed.

3 months when stored at -20°C to -80°C after reconstituting as directed.

THIS PRODUCT IS FOR RESEARCH USE ONLY AND IS NOT FOR USE IN HUMANS!

X

Quality Assurance