



Product Data Sheet

Recombinant Human BAFF / BLyS

Catalog Number: 100-24

Product Description

Alternate Names: TALL, BLyS, THANK

Accession Number: Q9Y275

Description: B cell-activating factor (BAFF), or B lymphocyte stimulator (BLyS), is a type II member of the tumor necrosis factor (TNF) superfamily. BAFF is expressed as a transmembrane protein on T cells, macrophages, and dendritic cells. The transmembrane domain of BAFF can also be cleaved to produce a soluble protein fragment. BAFF binds to the TNF receptors known as B cell maturation antigen (BCMA), transmembrane activator and CAML interactor (TACI), and BAFF receptor (BAFFR). BAFF is important for the survival and maturation of peripheral B cells. Human BAFF shows activity on mouse splenocytes.

Source: Genetically modified *E.coli*.

Predicted MW: Monomer, 18.5 kDa (with 163 amino acids)

AA Sequence: MHHHHHHLVP RAVQGPEETV TQDCLQLIAD SETPTIQKGS YTFVPWLLSF KRGSAL EEKE NKILVKETGY FFIYGQVLYT DKTYAMGHLI QRKKVHVFGD ELSLVTLFRC IQNMPETLPN NSCYSAGIAK LEEGDELQLA IPRENAQISL DGDVTFFGAL KLL

Formulation: Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5

Product Specifications*

*Lot-specific values for the following specifications are supplied with each production lot on its corresponding COA.

Specification	Method of Determination	Acceptance Criteria
Purity	Reducing and Non-Reducing SDS-PAGE	≥ 90%
Endotoxin	Kinetic LAL (50% confidence)	≤ 1 EUs/μg

Product Preparation and Storage

Country of Origin: USA

Shipping: Room temperature

Storage Prior to Reconstitution: -20°C

Product Reconstitution: Sterile water at 0.1 mg/mL

Instructions: Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws.

Expiration Date:

12 months from date of receipt when stored at -20°C to -80°C as supplied.

1 month when stored at 4°C after reconstituting as directed.

3 months when stored at -20°C to -80°C after reconstituting as directed.

THIS PRODUCT IS FOR RESEARCH USE ONLY AND IS NOT FOR USE IN HUMANS!