



Product Data Sheet

Recombinant Human RANK Ligand

Catalog Number: 100-19

Product Description

Alternate Names: TNFSF11, TRANCE, OPGL, ODF

Accession Number: O14788

Description: Receptor activator of nuclear factor kappa-B Ligand (RANK Ligand) is a cell-bound marker related to the tumor necrosis factor (TNF) family of proteins. RANK Ligand plays a critical role in bone metabolism and osteoclast differentiation. T cell expression of RANK Ligand promotes dendritic cell maturation.

Source: Genetically modified *E.coli*.

Predicted MW: Monomer, 19.7 kDa (with 175 amino acids)

AA Sequence: EKAMVDGSWL DLAKRSKLEA QPFAHLTINA TDIPSGSHKV SLSSWYHDRG WAKISNMTFS NGKLIVNQDG FYYLYANICF RHHETSGDLA TEYLQLMVYV TKTSIKIPSS HTLMKGGSTK YWSGNSEFHF YSINVGFFK LRSGEESIE VSNPSLLDPD QDATYFGAFK VRDID

Formulation: Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5

Product Specifications*

*Lot-specific values for the following specifications are supplied with each production lot on its corresponding COA.

Specification	Method of Determination	Acceptance Criteria
Purity	Reducing and Non-Reducing SDS-PAGE	≥ 95%
Endotoxin	Kinetic LAL (50% confidence)	≤ 1 EUs/μg
Biological Activity	Activity of RAW-Blue™ cells	ED50 ≤ 50 ng/mL (≥ 2.0 x 10 ⁴ units/mg)

In addition to the acceptance criteria, this product has been tested with the following assay:

(1) IL-8 production in human PBMCs

Product Preparation and Storage

Country of Origin: USA

Shipping: Room temperature

Storage Prior to Reconstitution: -20°C

Product Reconstitution: Sterile water at 0.1 mg/mL

Instructions: Centrifuge vial before opening. Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws.

Expiration Date:

12 months from date of receipt when stored at -20°C to -80°C as supplied.

1 month when stored at 4°C after reconstituting as directed.

3 months when stored at -20°C to -80°C after reconstituting as directed.

THIS PRODUCT IS FOR RESEARCH USE ONLY AND IS NOT FOR USE IN HUMANS!